



Help protect BC sturgeon

A survivor from the age of the dinosaurs over 200 million years ago, two ancient species of sturgeon found in BC waters today are experiencing population declines and are in need of protection: the **White Sturgeon**, which within Canada is unique to BC and is the largest freshwater fish in North America, and the **Green Sturgeon**, which is found along the west coast of North America.

Know how to identify these ancient species




Expect to find both sturgeon species in overlapping ranges in British Columbia.



Size, colour and age classes


While it can be tricky to differentiate the 2 species in younger life stages, a fully-grown, adult **White Sturgeon** that can live to be 100 years old, measures 3 times the length and 4 times the weight of an adult **Green Sturgeon**.



WHITE STURGEON

Acipenser transmontanus

Predominantly found in freshwater, White Sturgeon may also enter marine waters along the Pacific Coast between southern Alaska and Baja California. They are primarily observed in the Strait of Georgia and Fraser River estuary, and throughout the Strait of Juan de Fuca. **The Lower Fraser River population of White Sturgeon is currently assessed as Threatened** by the Committee on the Status of Endangered Wildlife in Canada (COSEWIC)



Scutes (bony plates)

White Sturgeon have prominent rows of scutes in front of the dorsal fin, along whitish lateral lines, and along the outsides of a clean, white belly.

Green Sturgeon are covered with smaller scutes and have 5 prominent rows of yellowish scutes with sharp tips on all sides of their bodies.



Barbels (whiskers)


White Sturgeon barbels are closer to the tip of the snout.

Green Sturgeon barbels are mid-way between the mouth and the tip of the snout, or closer to the mouth.

GREEN STURGEON

Acipenser medirostris



Predominantly found in saltwater along the BC coast, **Green Sturgeon** inhabit shallower areas of the continental shelf less than 110 meters deep and migrate into tidal areas of large coastal rivers and the same brackish estuaries as White Sturgeon. They are predominantly found along the coastlines of Washington, Oregon, Vancouver Island and Haida Gwaii. **Green Sturgeon is listed as a species of Special Concern** under Canada's *Species at Risk Act* (SARA).



Ventral features

White Sturgeon have 3-4 small pairs of pre-anal plates behind the pectoral fins and 1-2 post-anal plates behind the anal fin. The anal vent is located behind the pelvic fins.

Green Sturgeon pre- and post-anal plates vary in number, shape and distribution and the anal vent is located between the pelvic fins. A dark olive green stripe may or may not also be present on the belly in the upper torso region.



Sturgeon illustrations: Paul Vecsi; adapted from COSEWIC Status Report on the Green Sturgeon *Acipenser medirostris* in Canada 2004, Eddy and Underhill 1978



How to help BC sturgeon & their habitat

It is illegal to retain sturgeon in recreational and commercial fisheries.

Carefully and quickly release any incidental catch back into the water.

Protect sturgeon habitat by keeping it pollution-free.

Report any incidental sturgeon catch and fishing violations to Fisheries and Oceans Canada.

EXPECT to find White and Green Sturgeon in overlapping areas **PROTECT** sturgeon and their habitat **RESPECT** fishing regulations and closures